## Brandon M. Tauberg, M.D.

Orthopaedic Sports Medicine Surgeon Arthroscopic and Shoulder Surgery www.brandontaubergmd.com

## **Proximal Humerus Fracture Protocol**

Name:	l	Date:
Diagnosis:		
Date of Surgery/Injury:	_ Next Physician Appointment:	
Frequency: 1 2 3 4 times/week Duration	:1 2 3 4 5 6 Weeks	
<ul> <li>Weeks 0-2:</li> <li>Wear sling at all times except for exercise</li> <li>Elbow, wrist, hand ROM. Grip strengther</li> <li>Begin pendulum exercises</li> <li>Modalities: heat or ice</li> <li>Weeks 2-5:</li> <li>Wean from sling, d/c by 4-5 weeks</li> <li>Begin PROM exercises with gentle stretch</li> <li>Teach home exercise program</li> <li>Scapular retraction/depression</li> <li>Postural correction</li> <li>No resisted shoulder motion until 8 wee</li> <li>Modalities: Heat before PT, ice after PT,</li> <li>Weeks 5-12:</li> <li>Begin AROM, advance ROM as tolerated</li> <li>Light passive stretching at end ranges</li> <li>Begin scapular exercises, PRE's for large</li> <li>At 8 weeks, can begin strengthening/res</li> </ul>	ching at end ranges  eks post-op other modalities as needed  d to full  muscle groups (pecs, lats, etc)	
<ul> <li>Isometrics cuff strengthening with arm at Months 3-12 (if needed):</li> <li>Return to full motion</li> <li>Advance strengthening as tolerated: ison per rotator cuff, deltoid, and scapular st</li> <li>Only do strengthening 3x/week to avoid</li> <li>Maintenance home flexibility and exerci</li> <li>Sport or vocation specific rehabilitation</li> </ul>	metrics → bands → light weights (1-5 abilizers I rotator cuff tendonitis	5 lbs); 8-12 reps/2-3 sets
Comments:		
Teach Home Exercise Program		
Modalities: Heat before Ice after Other		Therapist's discretion
Signaturo:		